# **A Global Framework for Fertilizer BMPs**

By T.W. Bruulsema, C. Witt, Fernando García, Shutian Li, T. Nagendra Rao, Fang Chen, and S. Ivanova

This paper describes a framework designed to facilitate development and adoption of best management practices (BMPs) for fertilizer use, and to advance the understanding of how these practices contribute to the goals of sustainable development. The framework guides the application of scientific principles to determine which BMPs can be adapted to local conditions at the practical level.

t the farm level, cropping systems are managed for multiple objectives. Best management practices are those that most closely attain those objectives. Management of fertilizer use falls within a larger agronomic context of cropping system management. A framework is helpful for describing how BMPs for fertilizer use fit in with those for the agronomic system.

The goals of sustainable development, in the general sense, comprise equal emphasis on economic, social, and ecological aspects (Brundtland, 1987). Such development is essential to provide for the needs of current and future generations. At the farm level, however, it is difficult to relate specific crop management practices to these three general aspects. Four management objectives are applicable to the practical farm level of all cropping systems (Witt, 2003). These four objectives are productivity, profitability, cropping system sustainability, and a favorable biophysical and social environment (PPSE). They relate to each other as illustrated in **Figure 1**.

Fertilizer use BMPs comprise an interlinked subset of crop management BMPs. For a fertilizer use practice to be considered "best", it must harmonize with the other agronomic practices in providing an optimum combination of the four objectives, PPSE. It follows that the development, evaluation, and refinement of BMPs at the farm level must consider all four objectives, as must selection of indicators reflecting their combined impact at the regional, national, or global level. Appropriate indicators for use at different scales are further discussed below in the section on performance indicators.

# **Cropping System Management Objectives**


**Productivity.** For cropping systems, the primary measure of productivity is yield per unit area of cropland per unit of time. Productivity should be considered in terms of all resources, or production factors, involved. Several indicators describing production and input use efficiencies are probably required to properly evaluate productivity.

**Profitability.** Profitability is determined by the difference between the value of the produce (gross benefit or revenue) and the cost of production. Its primary measure is net benefit per unit of cropland per unit of time. The profitability gain of a specific management practice is the increase in gross revenue it generates, less its marginal cost.

Sustainability. Sustainability—at the level of the cropping system—refers to the influence of time on the resources involved. A sustainable production system is one in which the quality (or efficiency) of the resources used does not diminish over time, so that "outputs do not decrease when inputs are not increased" (Monteith, 1990).

Environment (biophysical and social). Crop production systems have a wide range of effects on surrounding





ecosystems through material losses to water and air. Specific effects can be limited to some extent by practices designed to optimize efficiency of resource use. Management choices at the farm level, when aggregated, also influence the social environment through demand for labor, working conditions, changes in ecosystem services, etc.

# **Fertilizer Management Objectives**

Fertilizer use BMPs essentially support the four objectives identified for cropping systems management and can be aptly described as the selection of the right source for application at the right rate, time, and place (Roberts, 2007). Fertilizer source, rate, timing, and placement are interdependent, and are also interlinked with the set of agronomic management practices applied in the cropping system, as illustrated in Figure 1.

# **Scientific Principles**

Specific scientific principles apply to crop and fertilizer use BMPs as a group and individually. These principles are both global and applicable at the practical farm management level. The application of these scientific principles may differ widely depending on the specific cropping system under consideration. **Specific principles relevant to each category of BMPs are listed below.** 

- 1) Crop Management
  - a) Seek practical measured validation.
  - b) Recognize and adapt to risks.
  - c) Define performance indicators.
  - d) Ensure two-way feedback between global and practical farm levels.
- 2) Fertilizer Management
  - a) Be consistent with understood process mechanisms.
  - b) Recognize interactions with other cropping system factors.
  - c) Recognize interactions among nutrient source, rate, time, and place.
  - d) Avoid detrimental effects on plant roots, leaves and seedlings.
  - e) Recognize effects on crop quality as well as yield.
  - f) Consider economics.
- 3) Source
  - a) Supply nutrients in plant-available forms.
  - b) Suit soil physical and chemical properties.
  - c) Recognize synergisms among nutrient elements and sources.
  - d) Recognize blend compatibility.
  - e) Recognize benefits and sensitivities to associated elements.
  - f) Control effects of non-nutritive elements.
- 4) Rate
  - a) Use adequate methods to assess soil nutrient supply.
  - b) Assess all indigenous nutrient sources available to the crop.
  - c) Assess crop demand for nutrients.
  - d) Predict fertilizer use efficiency.
  - e) Consider soil resource impacts.
  - f) Consider rate-specific economics.
- 5) Time
  - a) Assess timing of crop uptake.
  - b) Assess dynamics of soil nutrient supply.
  - c) Recognize timing of weather factors influencing nutrient loss.
  - d) Evaluate logistics of field operations.
- 6) Place
  - a) Recognize root-soil dynamics.
  - b) Manage spatial variability within fields and among farms.
  - c) Fit needs of tillage system.
  - d) Limit potential off-field transport of nutrients.

The number of scientific principles applicable to a given practical farming situation is considerable. Narrowing down to a set of BMPs appropriate to the practical level requires the involvement of qualified individuals: producers and advisers who understand both the principles and their application. Further details on these principles are provided in IPNI (2008).

# **Performance Indicators**

Performance indicators need to reflect the influence of fertilizer BMPs on all four crop management objectives. Nutrient use efficiency (NUE, yield or nutrient uptake per unit fertilizer nutrient applied) is often considered a foremost indicator relating to fertilizer use. However, as shown in Figure 1, it relates much more directly to profitability and productivity than it does to sustainability and environmental health. Other indicators of nutrient use efficiency exist (Dobermann, 2007; Snyder and Bruulsema, 2007) which differ in how well they relate to the four objectives. For example, one of the most important performance indicators for N is agronomic efficiency, the increase in grain yield per unit fertilizer nutrient applied. However, a low agronomic efficiency can be acceptable for nutrients such as P and K, for which a different measure of efficiency - partial nutrient balance - can be more relevant to the avoidance of soil nutrient depletion or excessive buildup.

The partial list of indicators shown in **Figure 1** is described further in **Table 1**. The set of performance indicators that describes the full impact of a combination of fertilizer BMPs varies depending on the scale of consideration. All stakeholders need to contribute to the selection of indicators for



optimum attainment of the four management objectives, PPSE. The framework concept we propose is helpful in ensuring that the set of indicators chosen provides a balanced reflection of the four objectives, in harmony with sustainable development goals.

# Conclusion

Best management practices for fertilizer use are those that support the achievement of the four main objectives of cropping systems management: productivity, profitability, sustainability, and environmental health. A strong set of scientific principles guiding the development and implementation of fertilizer use BMPs has evolved from a long history of agronomic and soil fertility research. Those principles—when seen as part of the global framework—show that the most appropriate set of fertilizer use BMPs can only be identified at the local level where the full context of each practice is known. The global framework for these BMPs also shows the need for employing a balanced complement of indicators to accurately describe the benefits and risks of fertilizer use in the context of sustainable development.

Dr. Bruulsema, Dr. Witt, Dr. García, Dr. Li, Dr. Rao, Dr. Chen, and Dr. Ivanova are IPNI scientific staff located in various regions of the world. All are members of the BMP Working Group of IPNI. Contact: Tom.Bruulsema@ipni.net.

Table 1. Performance indicators for fertilizer BMPs related to crop management objectives.		
Management Objective	Performance Indicator	Description
Productivity	Yield	Amount of crop harvested per unit of cropland per unit of time.
	Quality	Amounts of crop components harvested (sugar, protein, minerals, etc.) or other attributes that add value to the harvested product.
	Nutrient Use Efficiency	Yield or nutrient uptake per unit of nutrient applied.
	Water Use Efficiency	Yield per unit of water applied or available. Relevant to irrigated and rainfed production.
	Labor Use Efficiency	Labor demand and supply are critically linked to number and timing of field operations.
	Energy Use Efficiency	Crop yield per unit of energy input.
Profitability	Net Profit	Reflects both volume and value of crop produced, per unit of time, relative to all costs of production. Limitation is inability to deal with externalities that have not been attributed an economic value.
Cropping System Sustainability	Return on Investment	Similar to net profit, adding consideration of capital investment and amortization.
	Adoption	Proportion of producers using particular BMPs. Often easily measured, but context is important.
	Soil Productivity	Reflects changes in soil fertility levels, soil organic matter, and other soil quality indicators.
	Yield Stability	Resilience of crop yields to variations in weather and pests.
	Farm Income	Improvements in livelihood.
	Working conditions	Quality of life issues.
Healthy Social and Biophysical Environment	Water & Air Quality	Concentration and nutrient loading in water bodies of the agricultural watershed or airshed. Limited ability to monitor at farm scale; monitoring at the watershed, regional and global scales is an important public service.
	Ecosystem Services	Difficult to quantify. Important to identify. Can include crop dependence on natural predators and pollinators, link to outdoor recreation, hunting, fishing, etc.
	Biodiversity	Difficult to quantify - can be descriptive.
	Soil Erosion	Degree of soil coverage by actively growing crops and crop residues.
	Nutrient Loss	Specific losses of nutrients to water and air. Since there are many pathways, these can be difficult to measure at the farm level.
	Nutrient Balance	A total account of nutrient inputs and outputs, at the soil surface or farm gate. The requirement for nutrient inputs is often linked to the increasing nutrient removal with harvested products as yields increase.

# Acknowledgment

Dr. Paul Fixen contributed the groundwork for the framework concept, and his input through the process of its development is gratefully appreciated.

For more on this topic, visit the IPNI website at: >www.ipni.net/ conceptpapers<.



#### References

Brundtland, G.H. 1987. Our common future. Report of the World Commission on Environment and Development.

- Dobermann, A. 2007. Nutrient use efficiency measurement and management. pp 1-28. In Fertilizer Best Management Practices. IFA International Workshop on Fertilizer Best Management Practices (FBMPs). 7-9 March, 2007. Brussels, Belgium.
- IPNI. 2008. A global framework for best management practices for fertilizer use. IPNI Concept Paper #1. Norcross, GA.
- Monteith, J.L. 1990. Can sustainability be quantified? Indian J. Dryland Agric. Res. Dev. 5:1-5.
- Roberts, T.L. 2007. Right product, right rate, right time, and right place...the foundation of best management practices for fertilizer. pp. 29-32. In Fertilizer Best Management Practices. IFA International Workshop on Fertilizer Best Management Practices (FBMPs). 7-9 March, 2007. Brussels, Belgium.
- Snyder, C.S. and T.W. Bruulsema. 2007. Nutrient Use Efficiency and Effectiveness in North America: Indices of Agronomic and Environmental Benefit. International Plant Nutrition Institute. Reference # 07076.
- Witt, C. 2003. Fertilizer use efficiencies in irrigated rice in Asia. Proceedings of the IFA Regional Conference for Asia and the Pacific, Cheju Island, Republic of Korea, 6-8 October 2003. [online]. Available at www.fertilizer.org (last update 2003; accessed 27 Sept. 2005). Paris: International Fertilizer Association.